

Global Simulation of Sulfate Aerosol Using a Coupled Chemistry/Climate Model



C. R. Molenkamp, J. E. Penner, C. C. Chuang, J. J. Walton, C. S. Atherton, and C. J. O'Connor (All at Global Climate Research Division, Lawrence Livermore National Laboratory, Livermore, CA 94550; 510-422-1827; e-mail: cmole@llnl.gov)

C. Liousse (Centre des Faibles Radioactivités, CNRS-CEA, 91198 Gif sur Yvette, France; (1) 69 82 35 48)

Sulfate aerosols can affect global climate both directly by scattering solar radiation and indirectly by altering the cloud droplet size distribution. Therefore, knowledge of the spatial distribution of natural and anthropogenic sulfate aerosol is important.

We have recently coupled GRANTOUR, our global chemistry and transport model with the ECHAM climate model which provides several enhanced capabilities in the representation of aerosol interactions. ECHAM includes a specific representation of liquid water in large-scale clouds which allows us to represent wet phase gas-to-particle conversion of SO₂ to sulfate as well as improving the parameterization of precipitation scavenging.

Results from simulations will be compared to observations of scavenging and deposition of sulfate aerosols and to spatial distributions from earlier simulations based on CCM1. We also calculate the climate forcing due to anthropogenic sulfate as the difference in top of the atmosphere solar radiation when the solar radiation routine is called with and without the anthropogenic sulfate component.

* This work was performed under the auspices of the U. S. Department of Energy by the Lawrence Livermore National Laboratory under contract No. W-7405-ENG-48.

1. 1995 IUGG Meeting
2. NA
3. (a) Charles R. Molenkamp
L-262
Lawrence Livermore National
Laboratory
Livermore, CA 94550

(b) Tel: 1-510-422-1827

(c) Fax: 1-510-422-5844

(d) Email: cmole@llnl.gov
4. IAMAS
5. M W7 International Global
Aerosol Program
6. Poster = P
7. NA
8. No